

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com


CCR2 Ab

References(2) Images(8)

Cat.#: DF7507 Size:	Concn.: ~1mg/ml Source: Rabbit	Mol.Wt.: 42 kDa Clonality: Polyclonal
Application: Reactivity:	WB 1:1000-3000, IHC 1:50-1:200, IF/ICC 1:50-1:200 *The optimal dilutions should be determined by the end user. Human,Mouse,Rat	
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide a SulfoLink TM Coupling Resin (Thermo F	
Immunogen:	A synthesized peptide derived from hur within the internal amino acids.	nan CCR2, corresponding to a region
Uniprot:	P41597	



Western blot analysis of extracts from HepG2, using CCR2 Ab. Lane 1 was treated with the blocking peptide.



CCR2 at 1/100 staining Rat spinalis tissue sections by IF.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.