

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com


## FATP1 Ab

References(2) Images(5)

Cat.#: DF7716 Size:	Concn.: ~1mg/ml Source: Rabbit	Mol.Wt.: 65 kDa Clonality: Polyclonal
Application:	IF/ICC 1:100-1:500, WB 1:1000-3000 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human,Mouse,Rat	
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>™</sup> Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from hur region within the internal amino acids.	nan FATP1, corresponding to a
Uniprot:	Q6PCB7	



Western blot analysis of extracts from Myeloma cells, using FATP1 Ab. The lane on the left was treated with blocking peptide.



DF7716 staining Hela cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(#DF7716) and mouse anti-beta tubulin Ab(#T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI (blue).

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.