Phospho-CD45 (Ser1007) Ab

Images(3)

Cat.#: AF3127 Concn.: ~1mg/ml Mol.Wt.: 150kDa Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500

*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The Ab is from purified rabbit serum by affinity purification via sequential

chromatography on phospho-peptide and non-phospho-peptide affinity

columns.

Immunogen: A synthesized peptide derived from human CD45 around the


phosphorylation site of Ser1007.

Uniprot: P08575

Description: The protein encoded by this gene is a member of the protein tyrosine

phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation,


mitotic cycle, and oncogenic transformation.



Western blot analysis of extracts from Mouse testis, using Phospho-CD45

(Ser1007)[Ser1009] Ab at 1/1000 dilution.

Observed bands:125kD.



AF3127 staining HeLa cells by ICC/IF. Cells were fixed with PFA and permeabilized in 0.1% saponin prior to blocking in 10% serum for 45 minutes at 37°C. The primary Ab was diluted 1/400 and incubated with the sample for 1 hour at 37°C. A Alexa Fluor® 594 conjugated goat polyclonal to rabbit IgG (H+L), diluted 1/600 was used as secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4%C with gentle shaking,



overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.