

Phospho-Vitamin D Receptor (Ser208) Ab

[Images\(2\)](#)

Cat.#: AF3159 Concn.: ~1mg/ml Mol.Wt.: 45-55kD
 Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500
 *The optimal dilutions should be determined by the end user.

Reactivity: Human


Storage: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.

Purification: The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho-peptide and non-phospho-peptide affinity columns.


Immunogen: A synthesized peptide derived from human Vitamin D Receptor around the phosphorylation site of Ser208.

Uniprot: P11473

Description: Nuclear hormone receptor. VDR mediates the action of vitamin D3 by controlling the expression of hormone sensitive genes.



Western blot analysis of Phospho-Vitamin D Receptor (Ser208) Ab expression in heatshock treated HT29 cells lysates. The lane on the right was treated with the antigen-specific peptide.



AF3159 staining A549 cells cells by ICC/IF. Cells were fixed with PFA and permeabilized in 0.1% saponin prior to blocking in 10% serum for 45 minutes at 37°C. The primary Ab was diluted 1/400 and incubated with the sample for 1 hour at 37°C. A Alexa Fluor® 594 conjugated goat polyclonal to rabbit IgG (H+L), diluted 1/600 was used as secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.



Affinity Biosciences

website: www.affbiotech.com

order: order@affbiotech.com

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.