

Phospho-GFAP (Ser38) Ab

Images(5)

Cat.#: AF3166 Mol.Wt.: 50kDa Concn.: ~1mg/ml Source: Rabbit Size: Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500

*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% Storage:

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The Ab is from purified rabbit serum by affinity purification via sequential

chromatography on phospho-peptide and non-phospho-peptide affinity

columns.

Immunogen: A synthesized peptide derived from human GFAP around the

phosphorylation site of Ser38.

Uniprot: P14136

Description: This gene encodes one of the major intermediate filament proteins of mature

> astrocytes. It is used as a marker to distinguish astrocytes from other glial cells during development. Mutations in this gene cause Alexander disease, a rare disorder of astrocytes in the central nervous system. An additional transcript variant has been described, but its full length sequence has not

been determined.



Western blot analysis of extracts from Rat brain, using Phospho-GFAP (Ser38) Ab at 1/1000 dilution.

Observed bands:45kD.



AF3166 at 1/100 staining mouse brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary Ab.



<code>TMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.