

RCC2 Ab

Images(3)

Cat.#: AF9180 Size:	Concn.: ~1mg/ml Source: Rabbit	Mol.Wt.: 56kDa Clonality: Polyclonal
Application:	WB 1:1000, IHC 1:50-1:200 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human, Mouse, Rat	
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink [™] Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from hur within C-terminal amino acids.	man RCC2, corresponding to a region
Uniprot:	Q9P258	



Western blot analysis of extracts from K562, using RCC2 Ab. The lane on the left was treated with blocking peptide.



AF9180 at 1/100 staining Human spleen tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.