

C-MYC-tag Ab

[References\(3\)](#) [Images\(2\)](#)

Cat.#: T0052
Size:

Concn.: ~1mg/ml
Source: Rabbit

Mol.Wt.:
Clonality: Polyclonal

Application:

WB 1:3000-1:10000, IF/ICC: 1:200

*The optimal dilutions should be determined by the end user.

Reactivity:

All

Storage:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.

Purification:

Affinity-chromatography.

Immunogen:

A synthetic peptide EQKLISEEDL coupled to KLH.

Description:

c-Myc-tag antibody is part of the Tag series of antibodies, the best quality in the research. The immunogen of c-Myc tag antibody is a synthetic peptide corresponding to residues 410-419 of the human p62 c-myc protein conjugated to KLH. C-Myc tag antibody is suitable for detecting the expression level of c-Myc or its fusion proteins where the c-Myc tag is terminal or internal.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.