## Phospho-ZAP70 (Tyr319) Ab

References(1) Images(11)

Cat.#: AF3312 Concn.: ~1mg/ml Mol.Wt.: 70kDa
Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500

\*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The Ab is from purified rabbit serum by affinity purification via sequential

chromatography on phospho-peptide and non-phospho-peptide affinity

columns.

Immunogen: A synthesized peptide derived from human ZAP70 around the


phosphorylation site of Tyr319.

Uniprot: P43403


Description: ZAP70 (human) a tyrosine kinase of the Syk family. Associates with the T-

cell antigen receptor zeta-chain after TCR stimulation. Phosphorylated by Src-family kinases following antigen receptor activation. Plays a role in

lymphocyte activation.



AF3312 at 1/200 staining Rat kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary Ab.



AF3312 staining Hela cells(serum starvation) by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AF3312 1:200) and mouse anti-beta tubulin Ab(T0023 1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI(blue).

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at  $4^{\circ}$ C with gentle shaking,



overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.