## Phospho-DNA-PK (Thr2647) Ab

References(1) Images(3)

Cat.#: AF3360 Concn.: ~1mg/ml Mol.Wt.: 470kDa Size: Source: Rabbit Clonality: Polyclonal

Application: IHC 1:50-1:200, IF/ICC 1:100-1:500, WB 1:500-1:2000

\*The optimal dilutions should be determined by the end user.

Reactivity: Human

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The Ab is from purified rabbit serum by affinity purification via sequential

chromatography on phospho-peptide and non-phospho-peptide affinity

columns.

Immunogen: A synthesized peptide derived from human DNA-PK around the

phosphorylation site of Thr2647.

Uniprot: P78527

Description: The PRKDC gene encodes the catalytic subunit of a nuclear DNA-


dependent serine/threonine protein kinase (DNA-PK). The second

component is the autoimmune antigen Ku (MIM 152690), which is encoded

by the G22P1 gene on chromosome 22q.



AF3360 at 1/100 staining human brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary Ab.



AF3360 staining HUVEC cells treated with serum 20% 30' by ICC/IF. Cells were fixed with PFA and permeabilized in 0.1% saponin prior to blocking in 10% serum for 45 minutes at 37°C. The primary Ab was diluted 1/400 and incubated with the sample for 1 hour at 37°C. A Alexa Fluor 594 conjugated goat polyclonal to rabbit IgG (H+L), diluted 1/600 was used as secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4%C with gentle shaking,



overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.