

Phospho-Semaphorin 4B (Ser825) Ab

Images(1)

Concn.: ~1mg/ml	Mol.Wt.: 105kDa
Source: Rabbit	Clonality: Polyclonal
WB 1:500-1:2000, IHC 1:50-1:200	
*The optimal dilutions should be determ	ined by the end user.
Human,Mouse,Rat	
Rabbit IgG in phosphate buffered saline sodium azide and 50% glycerol. Store at date of receipt.	*
The Ab is from purified rabbit serum by chromatography on phospho-peptide and columns.	• 1 1
A synthesized peptide derived from hum phosphorylation site of Ser825.	an Semaphorin 4B around the
Q9NPR2	
	Source: Rabbit WB 1:500-1:2000, IHC 1:50-1:200 *The optimal dilutions should be determ Human,Mouse,Rat Rabbit IgG in phosphate buffered saline sodium azide and 50% glycerol. Store at date of receipt. The Ab is from purified rabbit serum by chromatography on phospho-peptide and columns. A synthesized peptide derived from hum phosphorylation site of Ser825.

<code>IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.</code>

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.