Phospho-HDAC2 (Ser394) Ab

Images(3)

Cat.#: AF3470 Concn.: ~1mg/ml Mol.Wt.: 60kDa Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500

*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The Ab is from purified rabbit serum by affinity purification via sequential

chromatography on phospho-peptide and non-phospho-peptide affinity

columns.

Immunogen: A synthesized peptide derived from human HDAC2 around the

phosphorylation site of Ser394.

Uniprot: Q92769

Description: HDAC2 a transcriptional regulator of the histone deacetylase family,

subfamily 1. Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation plays a role in epigenetic repression and transcriptional

regulation, cell cycle progression and developmental events.




Western blot analysis of extracts from various samples, using Phospho-

HDAC2 (Ser394) Ab.

Lane 1: Mouse muscle lysates; Lane 2: Mouse lung lysates;

Lane 3: Mouse lung lysates treated with blocking peptide;



AF3470 staining A549 cells(H2O2 treatment) by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(#AF3470) and mouse anti-beta tubulin Ab(#T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI (blue).



in 5% $\mbox{w/v}$ milk , 1% TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.