PDGF Receptor beta Ab

Images(1)

Cat.#: AF6134 Concn.: ~1mg/ml Mol.Wt.: 125kDa Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000

*The optimal dilutions should be determined by the end user.

Reactivity: Human,Rat

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLinkTM Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human PDGF Receptor beta,

corresponding to a region within the internal amino acids.

Uniprot: P09619

Description: This gene encodes a cell surface tyrosine kinase receptor for members of the

platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived growth

factor receptor alpha and beta polypeptides.

kDa 1 2 250-150-100-75-50-37-25-

Western blot analysis of PDGFRb expression in HepG2 whole cell lysates, The

lane on the left was treated with the antigen-specific peptide.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.