

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com


HGS Ab

Images(2)

Cat.#: AF6861 Size:	Concn.: ~1mg/ml Source: Rabbit	Mol.Wt.: 85~100kD Clonality: Polyclonal
Application:	IF/ICC 1:100-1:500, WB 1:500-1:2000 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human,Mouse,Rat	
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide a SulfoLink™ Coupling Resin (Thermo I	, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
Immunogen:	A synthesized peptide derived from hur	nan HGS.
Uniprot:	O14964	



Western blot analysis of extracts from HepG2 cells(heat shock treatment), using HGS Ab. The lane on the left was treated with blocking peptide.



AF6861 staining A549 cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AF6861) and mouse anti-beta tubulin Ab(T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI (blue).

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.