

Affinity Biosciences website:www.affbiotech.com

Cyclin D1 Ab

References(48) Images(36)

Cat.#: AF0931 Concn.: ~1mg/ml Mol.Wt.: 33kDa
Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500

*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLinkTM Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human Cyclin D1, corresponding to a

region within C-terminal amino acids.

Uniprot: P24385

Description: The protein encoded by this gene belongs to the highly conserved cyclin

family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclins function as regulators

of CDK kinases. Different cyclins exhibit distinct expression and

degradation patterns which contribute to the temporal coordination of each

mitotic event



Western blot analysis of extracts from various samples, using Cyclin D1 Ab.

Lane 1: RAW264.7 cells, blocked with antigen-specific peptides,

Lane 2: RAW264.7 cells,

Lane 3: C6 cells.



AF0931 at 1/100 staining rat brain by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22° C. An HRP conjugated goat anti-rabbit Ab was used as the secondary Ab.



in 5% $\mbox{w/v}$ milk , 1% TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.