

Affinity Biosciences

website:www.affbiotech.com order:order@affbiotech.com

CDH8 Ab

Images(3)

Cat.#: DF3533 Concn.: ~1mg/ml Mol.Wt.: 90 KD
Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:1000, IHC 1:50-1:200, IF/ICC 1:100-1:500

*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLinkTM Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human CDH8, corresponding to a

region within the internal amino acids.

Uniprot: P55286




Western blot analysis of extracts from various samples, using CDH8 Ab.

Lane 1: Hepg2 cells(heat-shock treatment), blocked with antigen-specific

peptides.

Lane 2: Hepg2 cells(heat-shock treatment). Lane 3: Hela cells(heat-shock treatment).



DF3533 staining Hela cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(DF3533) and mouse anti-beta tubulin Ab(T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI(blue).

IMPORTANT: For western blot, incubate membrane with diluted primary Ab
in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking,
overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.