

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com


CNN2 Ab

References(1) Images(2)

| Cat.#: DF3870 Size: | Concn.: ~1mg/ml Source: Rabbit | Mol.Wt.: 33 KD Clonality: Polyclonal |
|------------------------|--|---|
| Application: | WB 1:500-1:1000, IF/ICC 1:100-1:500 *The optimal dilutions should be determined by the end user. | |
| Reactivity: | Human, Mouse | |
| Storage: | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt. | |
| Purification: | The antiserum was purified by peptide affinity chromatography using SulfoLink [™] Coupling Resin (Thermo Fisher Scientific). | |
| Immunogen: | A synthesized peptide derived from hur region within the internal amino acids. | nan CNN2, corresponding to a |
| Uniprot: | Q99439 | |



Western blot analysis of extracts from mouse brain, using CNN2 Ab.



DF3870 staining A549 cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(DF3870) and mouse anti-beta tubulin Ab(T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI (blue).

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.