

IL-8 Ab

[Images\(1\)](#)

| | | |
|---------------|-----------------|-----------------------|
| Cat.#: BF0238 | Concn.: ~1mg/ml | Mol.Wt.: 11.1kDa |
| Size: | Source: Mouse | Clonality: Monoclonal |

Application: ELISA 1:10000, WB 1:500-1:2000
 *The optimal dilutions should be determined by the end user.

Reactivity: Human

Storage: Mouse IgG1 in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.

Purification: Affinity-chromatography.

Immunogen: Purified recombinant fragment of human IL-8 expressed in E. Coli.

Uniprot: P10145

Description: Interleukin-8 (IL-8) is a member of the CXC chemokine family. This chemokine is one of the major mediators of the inflammatory response. This chemokine is secreted by several cell types. It functions as a chemoattractant, and is also a potent angiogenic factor. IL-8 is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. IL-8 and other ten members of the CXC chemokine gene family form a chemokine gene cluster in a region mapped to chromosome 4q.




Figure 1: Western blot analysis using IL8 mouse mAb against IL8 recombinant protein.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.