

HPRT Ab

[Images\(1\)](#)

Cat.#: BF0205	Concn.: ~1mg/ml	Mol.Wt.: 25kDa
Size:	Source: Mouse	Clonality: Monoclonal

Application:	ELISA 1:10000, WB 1:500-1:2000 *The optimal dilutions should be determined by the end user.
Reactivity:	Human
Storage:	Mouse IgG1 in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.
Purification:	Affinity-chromatography.
Immunogen:	Purified recombinant fragment of human HPRT expressed in E. Coli.
Uniprot:	P00492
Description:	The HPRT1 gene provides instructions for making an enzyme called hypoxanthine phosphoribosyltransferase 1. This enzyme allows cells to recycle purines, some of the building blocks of DNA and its chemical cousin RNA. The enzyme hypoxanthine-guanine phosphoribosyltransferase (E.C.2.4.2.8., HPRT) plays a crucial role in uric acid synthesis and purine metabolism. This enzyme catalyzes the conversion of hypoxanthine and guanine to inosine monophosphate (IMP) and guanosine monophosphate (GMP), respectively, and uses phosphoribosylpyrophosphate (PRPP) as a cosubstrate and as a source of energy. This pathway is also known as the purine salvage pathway because it allows cells to reuse purine compounds to build DNA and RNA.




Figure 1: Western blot analysis using HPTR mouse mAb against truncated HPRT recombinant protein.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.