

ICAM2 Ab

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

Images(1)

Cat.#: DF4131 Size:	Concn.: ~1mg/ml Source: Rabbit	Mol.Wt.: 30 KD Clonality: Polyclonal
Application:	WB 1:500-1:1000, IF/ICC 1:100-1:500 *The optimal dilutions should be determined by the end user. Human,Mouse	
Reactivity:		
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink [™] Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from hun region within the internal amino acids.	nan ICAM2, corresponding to a
Uniprot:	P13598	

kDa 250-150-150-75-50-37-25-20-15-

Western blot analysis of extracts from NIH-3T3 cells, using ICAM2 Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.