

TLK2 Ab

Images(5)

| Cat.#: DF4478 Size: | Concn.: ~1mg/ml Source: Rabbit | Mol.Wt.: 90 KD Clonality: Polyclonal |
|------------------------|--|---|
| Application: | WB 1:500-1:1000, IF/ICC 1:100-1:500 *The optimal dilutions should be determined by the end user. Human,Mouse | |
| Reactivity: | | |
| Storage: | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt. | |
| Purification: | The antiserum was purified by peptide a SulfoLink [™] Coupling Resin (Thermo I | |
| Immunogen: | A synthesized peptide derived from hur within the internal amino acids. | nan TLK2, corresponding to a region |
| Uniprot: | Q86UE8 | |



Western blot analysis of extracts from EC304 cells(heat-shock treatment), using TLK2 Ab at 1/1000 dilution. Observed bands:85kD.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.