

Cytochrome P450 19A1 Ab

[References\(1\)](#) [Images\(2\)](#)

Cat.#: AF5229 Concn.: ~1mg/ml Mol.Wt.: 55 kDa
Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
*The optimal dilutions should be determined by the end user.

Reactivity: Human,Mouse,Rat


Storage: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).


Immunogen: A synthesized peptide derived from human Cytochrome P450 19A1, corresponding to a region within the internal amino acids.

Uniprot: P11511

Description: Defects in CYP19A1 are a cause of aromatase excess syndrome (AEXS) [MIM:139300]; also known as familial gynecomastia. AEXS is characterized by an estrogen excess due to an increased aromatase activity. Defects in CYP19A1 are the cause of aromatase deficiency (AROD) [MIM:107910]. AROD is a rare disease in which fetal androgens are not converted into estrogens due to placental aromatase deficiency.



Western blot analysis of Cytochrome P450 19A1 Ab expression in Human placenta tissue lysates. The lane on the left was treated with the antigen-specific peptide.



AF5229 at 1/100 staining human Placenta tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab

in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking,
overnight.

For Research Use Only. Not for use in diagnostic and therapeutic
procedures. Not for resale without express authorization.