## Affinity Biosciences

website:www.affbiotech.com order:order@affbiotech.com

## **NEUROD1 Ab**

Images(1)

Cat.#: DF6281 Concn.: ~1mg/ml Mol.Wt.: 40kDa Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500

\*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLink<sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human NEUROD1, corresponding to a

region within N-terminal amino acids.

Uniprot: Q13562

Description: NeuroD is a member of the basic helix-loop-helix (bHLH) family of

transcription factors. These proteins function by forming heterodimers with E-proteins and binding to the canonical E-box sequence CANNTG (1,2). Neuronal activity results in CaMKII-mediated phosphorylation of NeuroD at Ser336, which is necessary for formation and growth of dendrites (3,4). NeuroD is also phosphorylated at Ser274 though the results are context dependent as phosphorylation by Erk stimulates NeuroD activity in pancreatic?-cells while phosphorylation by GSK-3? inhibits NeuroD in neurons . NeuroD is crucially important in both the pancreas and developing nervous system, and plays a large role in the development of the inner ear

and mammalian retina.



Western blot analysis of extracts from various samples, using NEUROD1 Ab.

Lane 1: Rat spleen lysates; Lane 2: Rat brin lysates;

Lane 3: Rat brin lysates treated with blocking peptide;

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

procedures. Not for resale without express authorization.