

DPYD Ab

[Images\(2\)](#)

Cat.#: DF6507	Concn.: ~1mg/ml	Mol.Wt.: 110kDa
Size:	Source: Rabbit	Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
*The optimal dilutions should be determined by the end user.

Reactivity: Human,Mouse,Rat


Storage: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human DPYD, corresponding to a region within the internal amino acids.

Uniprot: Q12882

Description: Dihydropyrimidine dehydrogenase (DPD, DPYD) catalyzes the initial and rate-limiting step in uracil and thymidine catabolism as well as catabolism of the chemotherapeutic drug 5-fluorouracil (5-FU) and its derivatives. DPYD deficiency, which results from mutations in the DPYD gene, causes errors in pyrimidine metabolism and potentially life-threatening side effects in cancer patients treated with 5-FU (reviewed in 1). As a result, ongoing work examines whether or how DPYD gene variation and protein expression can be used to predict 5-FU toxicity (1,2). Several genes that impart resistance to 5-FU were recently identified in human hepatocellular carcinoma (HCC). AEG-1, which is highly expressed in HCC, increases the expression of DPYD.



Western blot analysis of extracts from rat brain, using DPYD Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.



Affinity Biosciences
website: www.affbiotech.com
order: order@affbiotech.com

procedures. Not for resale without express authorization.