

Cytochrome P450 11A1 Ab

[References\(6\)](#) [Images\(7\)](#)

Cat.#: DF6569	Concn.: ~1mg/ml	Mol.Wt.: 60kDa
Size:	Source: Rabbit	Clonality: Polyclonal
Application:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human,Mouse,Rat,Pig	
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from human Cytochrome P450 11A1, corresponding to a region within C-terminal amino acids.	
Uniprot:	P05108	
Description:	The formation of biologically active steroid hormones is significantly influenced by the mitochondrial cytochrome P450 enzyme, cholesterol side-chain cleavage. CYP11A1 functions in catalyzing the first step of steroid biosynthesis under modulation of cAMP. It is also known as cytochrome P450C11A1, cytochrome P450scc and cytochrome P450, subfamily XIA. Expression of CYP11A1 has been a useful tool for monitoring the differentiation state of cells from various endocrine tissues. Identification of CYP11A1 expression by 13363-1-AP detected a ~45kd band in human placenta, with respect to the reported band at 49-52kd by Roby, et al and Durkee, et al.	

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.