


## IL-10R alpha Ab

References(2) Images(4)

Cat.#: DF6643 Size:	Concn.: ~1mg/ml Source: Rabbit	Mol.Wt.: 63kDa Clonality: Polyclonal
Application:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human,Mouse	
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>™</sup> Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from hum a region within C-terminal amino acids.	an IL-10R alpha, corresponding to
Uniprot:	Q13651	
Description:	The protein encoded by this gene is a rec protein is structurally related to interfero mediate the immunosuppressive signal o the synthesis of proinflammatory cytokin promote survival of progenitor myeloid o substrate-2/PI 3-kinase/AKT pathway. A tyrosine phosphorylation of JAK1 and T variants, one protein-coding and the othe found for this gene. [provided by RefSec	n receptors. It has been shown to of interleukin 10, and thus inhibits nes. This receptor is reported to cells through the insulin receptor activation of this receptor leads to YK2 kinases. Two transcript er not protein-coding, have been



Western blot analysis of extracts from MCF-7, using IL10RA Ab. The lane on the left was treated with the antigen-specific peptide.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.