


VEGFR3 Ab

[Images\(1\)](#)

Cat.#: BF0536	Concn.: ~1mg/ml	Mol.Wt.: 145kDa
Size:	Source: Mouse	Clonality: Monoclonal

Application:	ELISA 1:10000, WB 1:500-1:2000 *The optimal dilutions should be determined by the end user.
Reactivity:	Human
Storage:	Mouse IgG1 in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.
Purification:	Affinity-chromatography.
Immunogen:	Purified recombinant fragment of human FLT4 expressed in E. Coli.
Uniprot:	P35916
Description:	This gene encodes a tyrosine kinase receptor for vascular endothelial growth factors C and D. The protein is thought to be involved in lymphangiogenesis and maintenance of the lymphatic endothelium. Mutations in this gene cause hereditary lymphedema type IA. Tissue specificity: Placenta, lung, heart, and kidney, does not seem to be expressed in pancreas and brain. VEGFR-3 is induced in all endothelial cells (EC's) during early embryogenesis, and its expression eventually disappears from the vascular endothelial cells of adult tissues. VEGFR-3 is constitutively expressed in the adult lymphatic endothelium.



Western blot analysis using FLT4 mAb against FLT4(AA: 25-330)-hIGFc transfected HEK293 cell lysate

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.