

## Cytochrome P450 19A1 Ab

[References\(1\)](#) [Images\(1\)](#)

Cat.#: DF6884	Concn.: ~1mg/ml	Mol.Wt.: 58kDa
Size:	Source: Rabbit	Clonality: Polyclonal
Application:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human,Mouse,Rat	
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from human Cytochrome P450 19A1, corresponding to a region within the internal amino acids.	
Uniprot:	P11511	
Description:	CYP19A1(Cytochrome P450 19A1) is also named as ARO1(aromatase), CYAR, CYP19, estrogen synthase and belongs to the cytochrome P450 family. It is a terminal enzyme which transforms irreversibly androgens into estrogens and it is present in the endoplasmic reticulum of numerous tissues(PMID:16406261). The gene encodes a protein with the molecular weight between 46 kDa and 69 kDa(PMID:8129748) and the protein can be glycosylated, but the level of glycosylation does not seem to be essential for CYP19A1 activity(PMID:12606587). Brain aromatase may be neuroprotective by increasing the local estrogen levels in injured neurons so that genetic variation in the brain aromatase gene may modify the risk for AD(PMID:16767510). This antibody is specific to CYP19A1.	

**IMPORTANT:** For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.