


p50 Dynamitin Ab

[Images\(1\)](#)

Cat.#: DF6923 Size:	Concn.: ~1mg/ml Source: Rabbit	Mol.Wt.: 44kDa Clonality: Polyclonal
Application:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human,Mouse,Rat	
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from human p50 Dynamitin, corresponding to a region within C-terminal amino acids.	
Uniprot:	Q13561	
Description:	Dynamitin is a 50 kDa protein containing a calmodulin binding domain, a putative ATPase domain and MacMARCKS-binding domain. This protein is a part of the dynactin complex believed to link the dynactin complex to membrane compartments. Its functions are tightly associated with dynein motor protein, thus extend to vesicle trafficking and membrane integrity. Dynamitin was named so because its overexpression causes dynactin complex which contains 10 subunits, to disassemble. Its N terminal 58 amino acid is for MacMARCKS binding and residues 59-83 is responsible for calmodulin binding. This antibody is against the full length p50 dynamitin.	



Western blot analysis of MCF7 cell lysates, using DCTN2 Ab. The lane on the left was treated with the antigen-specific peptide.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.