

Di-Methyl-Histone H3 (Lys4)/H3K4me2 Ab

[Images\(2\)](#)

Cat.#: DF6934 Concn.: ~1mg/ml Mol.Wt.: 15kDa
Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:50-1:200, IP 1:50-1:200,
CHIP 1:50-1:200

*The optimal dilutions should be determined by the end user.

Reactivity: Human,Mouse


Storage: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02%
sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from
date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using
SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthetic methylated peptide derived from human Di-Methyl-Histone H3
around the methylation site of Lys4.

Uniprot: Q16695

Description: Modulation of chromatin structure plays an important role in the regulation
of transcription in eukaryotes. The nucleosome, made up of DNA wound
around eight core histone proteins (two each of H2A, H2B, H3, and H4), is
the primary building block of chromatin . The amino-terminal tails of core
histones undergo various post-translational modifications, including
acetylation, phosphorylation, methylation, and ubiquitination (2-5). These
modifications occur in response to various stimuli and have a direct effect
on the accessibility of chromatin to transcription factors and, therefore, gene
expression . In most species, histone H2B is primarily acetylated at Lys5,
12, 15, and 20 (4,7). Histone H3 is primarily acetylated at Lys9, 14, 18, 23,
27, and 56.



Western blot analysis of extracts from HeLa using H3K4me2 Ab. The lane on the left was treated with the antigen-specific peptide.



DF6934 at 1/100 staining Human brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.