Affinity Biosciences website:www.affbiotech.com

mGluR8 Ab

Images(1)

Mol.Wt.: 102kDa Cat.#: DF7121 Concn.: ~1mg/ml Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200

*The optimal dilutions should be determined by the end user.

Human, Mouse, Rat Reactivity:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% Storage:

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLink™ Coupling Resin (Thermo Fisher Scientific).


A synthesized peptide derived from human mGluR8, corresponding to a Immunogen:

region within C-terminal amino acids.

O00222 Uniprot:

Description: L-glutamate is the major excitatory neurotransmitter in the central nervous

> system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8.



Western blot analysis of extracts from mouse brain tissue, using GRM8Ab. The lane on the left was treated with the antigen-specific peptide.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.