Affinity Biosciences website:www.affbiotech.com

SSTR1 Ab

Images(1)

Cat.#: DF7205 Concn.: ~1mg/ml Mol.Wt.: 43kDa Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200

*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLinkTM Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human SSTR1, corresponding to a

region within C-terminal amino acids.


Uniprot: P30872

Description: Somatostatins are peptide hormones that regulate diverse cellular functions

such as neurotransmission, cell proliferation, and endocrine signaling as well as inhibiting the release of many hormones and other secretory proteins. Somatostatin has two active forms of 14 and 28 amino acids. The biological effects of somatostatins are mediated by a family of G-protein coupled somatostatin receptors that are expressed in a tissue-specific manner. The protein encoded by this gene is a member of the superfamily of somatostatin receptors having seven transmembrane segments. Somatostatin receptors form homodimers and heterodimers with other members of the

superfamily as well as with other G-protein coupled receptors and receptor

tyrosine kinases.



Western blot analysis of THP-1 cell lysates, using SSTR1 Ab. The lane on the left was treated with the antigen-specific peptide.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

procedures. Not for resale without express authorization.