


NR2C2 Ab

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

Images(1)

Cat.#: DF2219 Size:	Concn.: ~1mg/ml Source: Rabbit	Mol.Wt.: 65 kDa Clonality: Polyclonal
Application: Reactivity:	WB 1:500-1:2000 *The optimal dilutions should be determined by the end user. Human.Mouse.Rat	
Storage:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from human NR2C2, corresponding to a region within the internal amino acids.	
Uniprot:	P49116	
Description:	Orphan nuclear receptor that can act as a repressor or activator of transcription. An important repressor of nuclear recptor signaling pathways such as retinoic acid receptor, retinoid X, vitamin D3 receptor, thyroid hormone receptor and estrogen receptor pathways. May regulate gene expression during the late phase of spermatogenesis. Together with NR2C1, forms the core of the DRED (direct repeat erythroid-definitive) complex that represses embryonic and fetal globin transcription including that of GATA1. Binds to hormone response elements (HREs) consisting of two 5'-AGGTCA-3' half site direct repeat consensus sequences. Plays a fundamental role in early embryonic development and embryonic stem cells. Required for normal spermatogenesis and cerebellum development.	



Western blot analysis of NR2C2 expression in 293T cell lysates.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic



procedures. Not for resale without express authorization.