

RAG1 Ab

Images(1)

Cat.#: BF0038 Size:	Concn.: ~1mg/ml Source: Mouse	Mol.Wt.: 119kDa Clonality: Monoclonal
5120.	Source. Mouse	Cionanty. Monocional
Application:	ELISA 1:10000, WB 1:500-1:2000 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human	
Storage:	Mouse IgG1 in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	Affinity-chromatography.	
Immunogen:	Purified recombinant fragment of huma	n RAG1 expressed in E. Coli.
Uniprot:	P15918	
Description:	The protein encoded by this gene is invo immunoglobulin V-D-J recombination. recognition of the DNA substrate, but st also requires RAG2. Defects in this gen diseases.	The encoded protein is involved in table binding and cleavage activity
kDa 1		




Figure 1: Western blot analysis using RAG1 mAb against human RAG1 recombinant protein. (Expected MW is 31.6 kDa)

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.