

AAK1 Ab

[Images\(1\)](#)

| | | |
|---------------|--|-----------------------|
| Cat.#: AF0685 | Concn.: ~1mg/ml | Mol.Wt.: 94kDa |
| Size: | Source: Rabbit | Clonality: Polyclonal |
| Application: | WB 1:500-1:2000, IF/ICC 1:100-1:500 *The optimal dilutions should be determined by the end user. | |
| Reactivity: | Human,Mouse,Rat,Monkey | |
| Storage: | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt. | |
| Purification: | The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific). | |
| Immunogen: | A synthesized peptide derived from human AAK1, corresponding to a region within the internal amino acids. | |
| Uniprot: | Q2M2I8 | |
| Description: | AAK1 is an NAK kinase. Plays a regulatory role in clathrin-mediated endocytosis, including the regulation of fatty acid synthesis and cholesterol synthesis. | |

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.